


Journal Pre-proof



Brain-derived neurotrophic factor (BDNF) polymorphism may influence the efficacy of tACS to modulate neural oscillations

Justin Riddle, Trevor McPherson, Alana K. Atkins, Christopher Walker, Sangtae Ahn, Flavio Frohlich

PII: S1935-861X(20)30091-7

DOI: <https://doi.org/10.1016/j.brs.2020.04.012>

Reference: BRS 1721

To appear in: *Brain Stimulation*

Received Date: 10 March 2020

Revised Date: 13 April 2020

Accepted Date: 17 April 2020

Please cite this article as: Riddle J, McPherson T, Atkins AK, Walker C, Ahn S, Frohlich F, Brain-derived neurotrophic factor (BDNF) polymorphism may influence the efficacy of tACS to modulate neural oscillations, *Brain Stimulation*, <https://doi.org/10.1016/j.brs.2020.04.012>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier Inc.

Brain-derived neurotrophic factor (BDNF) polymorphism may influence the efficacy of tACS to modulate neural oscillations

Dear Editor,

Plasticity of human motor cortex is modulated by brain-derived neurotrophic factor (BDNF). A single nucleotide polymorphism at codon 66 (Val66Met) in the gene that codes for BDNF is associated with altered learning and memory [1]. Non-invasive brain stimulation increases excitability by recruiting BDNF-dependent neural plasticity [2]. In a previous study, the presence of the Val66Met polymorphism predicted reduced efficacy for both transcranial current stimulation and repetitive transcranial magnetic stimulation (TMS) to modulate excitability in the corticospinal tract probed as the amplitude of the motor-evoked potential elicited by TMS pulses to primary motor cortex [3]. In animal models, the Val66Met polymorphism reduces the neural response to brain stimulation by decreasing NMDA-dependent long-term potentiation of synaptic transmission [4, 5]. Despite this growing literature on the potential gating of corticospinal excitability changes by the Val66Met polymorphism, the role of the Val66Met polymorphism in gating the neural response to brain stimulation at the level of brain network dynamics remains unknown. Alpha oscillations are the dominant brain rhythm in the awake resting state and modulate the neural response to sensory input [6]. Given this role in shaping state-dependent neural excitability, targeting alpha oscillations with brain stimulation represents a promising strategy to modulate and enhance brain function. In particular, transcranial alternating current stimulation with stimulation waveforms in the alpha frequency (alpha-tACS) modulate alpha oscillations and were recently investigated as a treatment for major depressive disorder [7], schizophrenia [8], and chronic pain [9]. Yet, the question remains if the modulation of alpha oscillations by alpha-tACS is a function of Val66Met status. We analyzed EEG data from three alpha-tACS studies in our group as a function of Val66Met polymorphism.

All data were collected in double-blind, placebo-controlled studies performed at the University of North Carolina (UNC) at Chapel Hill. All study protocols were approved by the local Institutional Review Board at UNC and all study participants provided written informed consent. The data presented here spans from August 2017 to August 2019. The first study investigated the effect of tACS on alpha oscillations in healthy controls in a cross-over design (unpublished). The second study investigated the effect of tACS on alpha oscillations in patients with chronic back pain [9]. The third study investigated the effect of tACS in depression and included a set of healthy control participants that are included in the analysis here (unpublished). In all studies included in this paper, we obtained a saliva sample from the participants using the Oragene Discover OGR-500 DNA Self-Collection kits (Genotek Inc., Ottawa, ON, Canada) for subsequent genotyping. DNA was extracted and sequenced at the Val66Met locus by Genotek (Ottawa, Canada) using the TaqMan SNP Genotyping assay. EEG data were analyzed as previously described [9]. We defined an a priori area of interest centered on the central EEG electrode Cz (group of six electrodes), where we have previously found an increase in alpha oscillation with the electrode montage used in these studies (2mA peak-to-peak at bilateral frontal electrode and 4mA peak-to-peak at central posterior “return” electrode) [9]. The amplitude of the alpha oscillation was log transformed and spatially normalized by z-scoring.

In our pooled dataset, we found 58 Val/Val, 37 Val/Met, and 5 Met/Met participants. We combined the latter two groups into a single Met carrier group. In agreement with our hypothesis, we found that the increase of the amplitude of the alpha oscillation by alpha-tACS (verum versus placebo) was a function of the Val66Met polymorphism ($F(1,96) = 4.350$, $p = 0.0397$, $\eta_p^2 = 0.043$) by comparing the Val66Val ($N = 58$) with the Met carrier group ($N = 42$).

This finding points towards a role of BDNF-dependent plasticity in the offline, or consequent, effect of tACS. Intriguingly, a previous study concluded that plasticity explains the offline effect of tACS by demonstrating no significant difference in the effect of tACS when delivered in consecutive periods that were either phase-continuous or phase-discontinuous [10]. However, no genotyping was performed. To

our knowledge, we are the first to examine how the BDNF Val66Met polymorphism affects the frequency-specific response of neural oscillation to tACS.

As with any scientific investigation, our study has several limitations. First, we are unable to answer the question whether the online, or concurrent, effect of tACS depends on the Val66Met polymorphism. Such mechanistic insights of entrainment during tACS in human EEG studies would require the recording of neural activity during stimulation and successful removal of the stimulation artefact, which remains a technical challenge. Second, we pooled across studies where some used 10Hz-tACS (Studies 1 and 2) and one used tACS at the individual alpha frequency (Study 3). In addition, the position of the return electrode slightly differed as in that the location of the third electrode was more anterior (Cz) for Study 3 in comparison to Studies 1 and 2 (POz). However, the two in-phase frontal electrodes were located at F3 and F4 in all three studies. Finally, we have pooled data from three different studies where the primary outcome was not to test the role of BDNF. Thus, other unaccounted differences between the three studies may play a role in our results.

In summary, our results support further investigations of how the Val66Met polymorphism shapes the response of neural oscillations to tACS. Assessing this SNP may be of particular relevance in tACS treatment clinical trials with multiple stimulation sessions.

Sincerely,

Justin Riddle, Trevor McPherson, Alana K. Atkins, Christopher Walker, Sangtae Ahn, Flavio Frohlich

Acknowledgements

We would like to thank all the past and current members of the Frohlich Lab and the Carolina Center for Neurostimulation who have contributed to the studies included in this letter.

Conflict of Interest

F.F. is the lead inventor of IP filed by UNC. F.F. is the founder, CSO, and majority owner of Pulvinar Neuro LLC.

Funding

The work presented here was partially supported by the National Institute of Mental Health and the National Institutes of Health (award R01MH101547, to F.F.). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

REFERENCES

1. Kleim, J.A., et al., *BDNF val66met polymorphism is associated with modified experience-dependent plasticity in human motor cortex*. Nature Neuroscience, 2006. **9**(6): p. 735-737.
2. Cheeran, B., et al., *A common polymorphism in the brain-derived neurotrophic factor gene (BDNF) modulates human cortical plasticity and the response to rTMS*. The Journal of Physiology, 2008. **586**(23): p. 5717-5725.
3. Antal, A., et al., *Brain-derived neurotrophic factor (BDNF) gene polymorphisms shape cortical plasticity in humans*. Brain Stimulation, 2010. **3**(4): p. 230-237.
4. Fritsch, B., et al., *Direct Current Stimulation Promotes BDNF-Dependent Synaptic Plasticity: Potential Implications for Motor Learning*. Neuron, 2010. **66**(2): p. 198-204.
5. Podda, M.V., et al., *Anodal transcranial direct current stimulation boosts synaptic plasticity and memory in mice via epigenetic regulation of Bdnf expression*. Scientific Reports, 2016. **6**.
6. Klimesch, W., *Alpha-band oscillations, attention, and controlled access to stored information*. Trends in Cognitive Sciences, 2012. **16**(12): p. 606-617.
7. Alexander, M.L., et al., *Double-blind, randomized pilot clinical trial targeting alpha oscillations with transcranial alternating current stimulation (tACS) for the treatment of major depressive disorder (MDD)*. Translational Psychiatry, 2019. **9**(1): p. 106.
8. Ahn, S., et al., *Targeting reduced neural oscillations in patients with schizophrenia by transcranial alternating current stimulation*. NeuroImage, 2019. **186**: p. 126-136.
9. Ahn, S., et al., *Identifying and Engaging Neuronal Oscillations by Transcranial Alternating Current Stimulation in Patients With Chronic Low Back Pain: A Randomized, Crossover, Double-Blind, Sham-Controlled Pilot Study*. The Journal of Pain, 2018.
10. Vossen, A., J. Gross, and G. Thut, *Alpha power increase after transcranial alternating current stimulation at alpha frequency (α -tACS) reflects plastic changes rather than entrainment*. Brain stimulation, 2015. **8**(3): p. 499-508.